



Influenza vaccination during pregnancy (and in women planning pregnancy)

This statement has been developed and reviewed by the Women's Health Committee and approved by the RANZCOG Board and Council.

A list of Women's Health Committee Members can be found in [Appendix A](#).

Disclosure statements have been received from all members of this committee.

Disclaimer: This College Statement is intended to provide general advice to practitioners. The statement should never be relied on as a substitute for proper assessment with respect to the particular circumstances of each case and the needs of each patient. This document reflects emerging clinical and scientific advances as of the date issued and is subject to change. The statement has been prepared having regard to general circumstances.

First endorsed by RANZCOG: November 2011
Current: March 2017
Review due: March 2020

Objectives: To provide advice on influenza and vaccination during pregnancy and in women planning pregnancy.

Options: Immunisation against influenza versus no immunisation.

Outcomes: To minimise the incidence of influenza infection in pregnant women, in women planning pregnancy and in babies up to 6 months of age.

Target audience: All health practitioners providing maternity care and patients.

Validation: This statement is in line with the recommendations in the 10th edition of the Australian Immunisation Handbook.¹

Funding: The development and review of this statement was funded by RANZCOG.

Background: This statement was originally developed in November 2011 and reviewed in July 2013 following publication of the 10th edition of the Australian Immunisation Handbook¹. The latest review was undertaken in March 2017.

Table of Contents

1. Patient summary.....	3
2. Summary of recommendations	3
3. Introduction.....	4
4. Evidence Summary and Recommendations	4
4.1 What are the risks associated with influenza infection during pregnancy?	4
4.2 Should pregnant women or those planning pregnancy be vaccinated against influenza?	4
4.3 What is the optimal timing of influenza vaccination in pregnant women or those planning pregnancy?	5
4.4 Are there any adverse consequences to vaccinating pregnant women or those planning pregnancy against influenza?.....	6
4.5 Does influenza vaccination during pregnancy provide protection for infants?.....	6
4.6 How long after vaccination can protection be expected?.....	6
4.7 Should obstetric and midwifery staff be vaccinated against influenza?.....	6
5. Links to other College statements.....	6
6. Patient information.....	8
7. References	7
Appendices.....	8
Appendix A Women’s Health Committee Membership	9
Appendix B Overview of the development and review process for this statement	9
Appendix C Full Disclaimer.....	11

1. Patient summary

Pregnant women are at increased risk of severe consequences of influenza infection. This is because there are a number of changes that occur to a woman's body during pregnancy which may put pregnant women at higher risk of complications from the flu (e.g. changes to lung function, increased cardiac output, increased oxygen consumption, and changes to the immune response).

The best way to protect pregnant women against flu is by vaccinating against it. Vaccination needs to be given each year because the viruses are always changing. Vaccinating against flu during pregnancy provides protection for the mother and the new born baby for the first six months after birth.

The influenza vaccine is safe for pregnant women in all trimesters, with no unusual patterns in pregnancy or fetal outcomes being observed in vaccine adverse events reports.

2. Summary of recommendations

Recommendation 1	Grade and reference
Influenza vaccination is recommended for all pregnant women regardless of gestation, and in women planning pregnancy.	Consensus-based recommendation 1
Good Practice Note	Grade
Free influenza vaccine is available to all pregnant women in Australia and New Zealand . To receive the influenza vaccination, pregnant women are advised to visit their local doctor or immunisation provider. It is important to note that the vaccine is free; however a consultation fee may apply. For more information relevant to Australia please visit: http://www.health.gov.au/internet/immunise/publishing.nsf/Content/immunise-influenza or http://www.influenza.org.nz/?t=887 for information specific to New Zealand.	Good Practice Note
Recommendation 2	Grade
Vaccination early in the season and regardless of gestational age is optimal, but unvaccinated pregnant women should be immunised at any time during influenza season as long as the vaccine supply lasts.	Consensus-based recommendation
Recommendation 3	Grade
The Royal Australian and New Zealand College of Obstetricians and Gynaecologists strongly endorse routine vaccination of obstetric and midwifery staff, both to protect these individuals as well as their families, close contacts and patients.	Consensus-based recommendation

3. Introduction

Influenza is a very serious and potentially life-threatening illness, caused by the influenza virus. Major outbreaks associated with severe disease are usually caused influenza virus types A and B. While influenza B is a human disease, subtypes of influenza A, characterized by differing haemagglutinin (H) and neuraminidase (N) antigens, infect many species. The H and N antigens frequently change their protein structure, leading to the development of new strains of influenza A virus that may infect humans and cause serious disease. For this reason, new vaccination preparations must be developed for each pandemic season.

Although most healthy people will suffer a serious but self-limited illness lasting one to two weeks, some groups are particularly vulnerable to significant disease and potentially death. Pregnant women are more likely to become seriously ill and to die than the general population. This vulnerability probably relates to changes in maternal physiology, including changes to the respiratory and cardiovascular system with reduced lung capacity and changes in the immune system.

Influenza vaccination during pregnancy should be routine: safety is well established²⁻¹⁰ and both maternal and infant benefit is now proven with only 5 vaccination doses estimated to prevent one case of maternal or infant respiratory illness.⁹

4. Evidence Summary and Recommendations

4.1 What are the risks associated with influenza infection during pregnancy?

Preventing influenza during pregnancy is an essential part of antenatal care because pregnant women are at an increased risk of serious illness due to influenza.¹¹ Excess morbidity and mortality in pregnant women infected with influenza compared with non-pregnant women of similar age who are infected with influenza has been noted during pandemics as long ago as 1918¹² but drew public and professional attention most recently during 2009.^{13, 14}

The risk to the mother of complications from influenza increases in the later stages of pregnancy.^{2-5, 9, 15-19} Infection in the third trimester of pregnancy appears to be the most dangerous for the pregnant woman.¹⁴

4.2 Should pregnant women or those planning pregnancy be vaccinated against influenza?

The most effective strategy for preventing influenza in pregnant women is annual immunisation. Influenza vaccination is estimated to prevent 1 to 2 hospitalisations per 1000 women vaccinated during the second or third trimester.¹

The Australian Government Department of Health and Ageing strongly recommends vaccination during the influenza season for pregnant women and women planning pregnancy. ([see Chapter 4.7](#)).¹

The [New Zealand Ministry of Health](#) strongly recommends seasonal influenza vaccination for all women who will be pregnant during the influenza season (which is usually between May to October).

The Royal Australian and New Zealand College of Obstetricians and Gynaecologists strongly endorse routine influenza vaccination of pregnant women and women planning pregnancy.

In Australia and New Zealand, the flu season is considered to be between May to October, usually peaking by the end of August.²⁰

The influenza vaccine is available for free to all pregnant women in [Australia](#) and [New Zealand](#).

Recommendation 1	Grade and reference
Influenza vaccination is recommended for all pregnant women regardless of gestation, and for women planning pregnancy.	Consensus-based recommendation
Good Practice Note	Grade
<p>Free influenza vaccine is available to all pregnant women in Australia and New Zealand.</p> <p>To receive the influenza vaccination, pregnant women are advised to visit their local doctor or immunisation provider. It is important to note that the vaccine is free; however a consultation fee may apply.</p> <p>For more information relevant to Australia please visit: http://www.health.gov.au/internet/immunise/publishing.nsf/Content/immunise-influenza or http://www.influenza.org.nz/?t=887 for information specific to New Zealand.</p>	Good Practice Notes

4.3 What is the optimal timing of influenza vaccination in pregnant women or those planning pregnancy?

Inactivated influenza vaccine is usually available from February each year in the Southern Hemisphere. Live attenuated influenza vaccination has not been licensed in Australia.¹

Vaccination early in the season and regardless of gestational age is optimal, but unvaccinated pregnant women should be immunised at any time during influenza season as long as the vaccine supply lasts.

In Australia and New Zealand, the flu season is considered to be between May to October, usually peaking by the end of August.²⁰

Whilst it is recommended that all pregnant women should be immunised as early as possible in pregnancy,²¹ the precise timing of vaccination will depend on the time of year, vaccine availability, influenza seasonality, gestation of pregnancy and the likely duration of immunity.

The vaccine should be administered annually to maintain immunity and to provide protection against new strains.

Recommendation 2	Grade
Vaccination early in the season and regardless of gestational age is optimal, but unvaccinated pregnant women should be immunised at any time during influenza season as long as the vaccine supply lasts.	Consensus-based recommendation

4.4 Are there any adverse consequences to vaccinating pregnant women or those planning pregnancy against influenza?

No study to date has shown an adverse consequence of inactivated influenza vaccine in pregnant women or their offspring.^{2-5, 7-9 18,22, 23}

Influenza vaccination is safe at any stage of pregnancy and there is no preference for any specific type of influenza vaccine in pregnancy; all are considered acceptable for use in pregnant women.

The only absolute contraindications to influenza vaccines are:

- anaphylaxis following a previous dose of any influenza vaccine; or
- anaphylaxis following any vaccine component.

See section 4.7.10 of the 10th edition Australian Immunisation Handbook for precautions for persons with a known egg allergy.¹

4.5 Does influenza vaccination during pregnancy provide protection for infants?

Active placental transfer of maternal antibodies makes influenza vaccine during pregnancy a highly effective measure to protect infants from influenza during the first 6 months of life.^{9, 15, 24, 25} Most evidence around infant protection is from studies of maternal influenza vaccination in the second or third trimester.^{2-5, 9, 15-19, 24} No study to date has shown an adverse consequence of inactivated influenza vaccine in pregnant women or their offspring.^{6, 10, 19, 26}

4.6 How long after vaccination can protection be expected?

Some maternal benefit is might accrue as early as 2 weeks after vaccination with research in pregnant women demonstrating seroconversion by 4 to 6 weeks after vaccination.⁵

4.7 Should obstetric and midwifery staff be vaccinated against influenza?

Recommendation 3	Grade and reference
The Royal Australian and New Zealand College of Obstetricians and Gynaecologists strongly endorse routine vaccination of obstetric and midwifery staff, both to protect these individuals as well as their families, closes contacts and patients.	Consensus-based recommendation

5. References

1. Australian Government. The Australian Immunisation Handbook: 10th Edition 2013 Canberra2013. Available from: <http://www.health.gov.au/internet/immunise/publishing.nsf/Content/Handbook10-home>.
2. American College of Obstetricians and Gynecologists (ACOG). Committee Opinion No. 468 Influenza vaccination during pregnancy. *Obstetrics and gynecology*. 2010;116(10):1006–7.
3. Black SB, Shinefield HR, France EK, Fireman BH, Platt ST, Shay D, et al. Effectiveness of influenza vaccine during pregnancy in preventing hospitalizations and outpatient visits for respiratory illness in pregnant women and their infants. *American journal of perinatology*. 2004;21(6):333-9.
4. Irving WL, James DK, Stephenson T, Laing P, Jameson C, Oxford JS, et al. Influenza virus infection in the second and third trimesters of pregnancy: a clinical and seroepidemiological study. *BJOG : an international journal of obstetrics and gynaecology*. 2000;107(10):1282-9.
5. Mak TK, Mangtani P, Leese J, Watson JM, Pfeifer D. Influenza vaccination in pregnancy: current evidence and selected national policies. *The Lancet infectious diseases*. 2008;8(1):44-52.
6. Moro PL, Broder K, Zheteyeva Y, Walton K, Rohan P, Sutherland A, et al. Adverse events in pregnant women following administration of trivalent inactivated influenza vaccine and live attenuated influenza vaccine in the Vaccine Adverse Event Reporting System, 1990-2009. *American journal of obstetrics and gynecology*. 2011;204(2):146 e1-7.
7. Sheffield JS, Greer LG, Rogers VL, Roberts SW, Lytle H, McIntire DD, et al. Effect of influenza vaccination in the first trimester of pregnancy. *Obstetrics and gynecology*. 2012;120(3):532-7.
8. World Health Organization. Global Advisory Committee on Vaccine Safety, June 2012. *Weekly Epidemiological Record*. 2012(87):281-7.
9. Zaman K, Roy E, Arifeen SE, Rahman M, Raqib R, Wilson E, et al. Effectiveness of maternal influenza immunization in mothers and infants. *The New England journal of medicine*. 2008;359(15):1555-64.
10. Moro P BJ, Lewis P, Cragan J, Tepper N, Cano M., Surveillance of Adverse Events After Seasonal Influenza Vaccination in Pregnant Women and Their Infants in the Vaccine Adverse Event Reporting System, July 2010-May 2016. *Drug Safety*. 2017;40(2):145-52.
11. Rothberg MB, Haessler SD, Brown RB. Complications of viral influenza. *The American journal of medicine*. 2008;121(4):258-64.
12. Naleway AL, Smith WJ, Mullooly JP. Delivering influenza vaccine to pregnant women. *Epidemiologic reviews*. 2006;28:47-53.
13. Investigators AI, Webb SA, Pettila V, Seppelt I, Bellomo R, Bailey M, et al. Critical care services and 2009 H1N1 influenza in Australia and New Zealand. *The New England journal of medicine*. 2009;361(20):1925-34.
14. Hewagama S, Walker SP, Stuart RL, Gordon C, Johnson PD, Friedman ND, et al. 2009 H1N1 influenza A and pregnancy outcomes in Victoria, Australia. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America*. 2010;50(5):686-90.
15. Eick AA, Uyeki TM, Klimov A, Hall H, Reid R, Santosham M, et al. Maternal influenza vaccination and effect on influenza virus infection in young infants. *Archives of pediatrics & adolescent medicine*. 2011;165(2):104-11.
16. Englund JA, Mbawuikie IN, Hammill H, Holleman MC, Baxter BD, Glezen WP. Maternal immunization with influenza or tetanus toxoid vaccine for passive antibody protection in young infants. *The Journal of infectious diseases*. 1993;168(3):647-56.
17. France EK, Smith-Ray R, McClure D, Hambidge S, Xu S, Yamasaki K, et al. Impact of maternal influenza vaccination during pregnancy on the incidence of acute respiratory illness visits among infants. *Archives of pediatrics & adolescent medicine*. 2006;160(12):1277-83.

18. MacDonald NE, Riley LE, Steinhoff MC. Influenza immunization in pregnancy. *Obstetrics and gynecology*. 2009;114(2 Pt 1):365-8.
19. Tamma PD, Steinhoff MC, Omer SB. Influenza infection and vaccination in pregnant women. *Expert review of respiratory medicine*. 2010;4(3):321-8.
20. Australian Government: Department of Health. Australian influenza report 2013— 14 September to 27 September 2013 (#08/2013) 2013 [24 October 2013]. Available from: <http://www.health.gov.au/internet/main/publishing.nsf/content/cda-surveil-ozflu-flucurr.htm>.
21. Omer SB, Goodman D, Steinhoff MC, Rochat R, Klugman KP, Stoll BJ, et al. Maternal influenza immunization and reduced likelihood of prematurity and small for gestational age births: a retrospective cohort study. *PLoS medicine*. 2011;8(5):e1000441.
22. Munoz FM. Safety of influenza vaccines in pregnant women. *American journal of obstetrics and gynecology*. 2012;207(3):S33-S7.
23. Regan A, Moore, H., deKlerk, N., Omer, S., Shellam, G., Mak, D., Effler, P.,. Seasonal Trivalent Influenza Vaccination During Pregnancy and the Incidence of Stillbirth: Population-Based Retrospective Cohort Study. *Clinical Infectious Diseases: an official publication of the Infectious Diseases Society of America*. 2016;62(10):1221-7.
24. Benowitz I, Esposito DB, Gracey KD, Shapiro ED, Vazquez M. Influenza vaccine given to pregnant women reduces hospitalization due to influenza in their infants. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America*. 2010;51(12):1355-61.
25. Poehling KA, Szilagyi PG, Staat MA, Snively BM, Payne DC, Bridges CB, et al. Impact of maternal immunization on influenza hospitalizations in infants. *American journal of obstetrics and gynecology*. 2011;204(6 Suppl 1):S141-8.
26. Lacroix I, Damase-Michel C, Kreft-Jais C, Castot A, Montastruc JL, French Association of Regional Pharmacovigilance C. 2009 H1N1 influenza vaccines in pregnant women: the French Pharmacovigilance survey. *Vaccine*. 2011;29(7):1357-8.

6. Links to other College statements

Evidence-based Medicine, Obstetrics and Gynaecology (C-Gen 15)

http://www.ranzcog.edu.au/component/docman/doc_download/894-c-gen-15-evidence-based-medicine-obstetrics-and-gynaecology.html?Itemid=341

7. Patient information

NZ patient information - [Pregnant Women, Influenza and Influenza Vaccination and Frequently Asked Questions](#)

Australian Department of Health - [Influenza \(Flu\)](#)

Appendices

Appendix A Women's Health Committee Membership

Name	Position on Committee
Professor Yee Leung	Chair
Dr Joseph Sgroi	Deputy Chair, Gynaecology
Associate Professor Janet Vaughan	Deputy Chair, Obstetrics
Professor Susan Walker	Member
Associate Professor Ian Pettigrew	Member
Dr Tal Jacobson	Member
Dr Ian Page	Member
Dr John Regan	Member
Dr Craig Skidmore	Member
Dr Lisa Hui	Member
Dr Bernadette White	Member
Dr Scott White	Member
Associate Professor Kirsten Black	Member
Dr Greg Fox	College Medical Officer
Dr Marilyn Clarke	Chair of the A&TSI WHC
Dr Martin Byrne	GPOAC Representative
Ms Catherine Whitby	Community Representative
Ms Sherryn Elworthy	Midwifery Representative
Dr Amelia Ryan	Trainee Representative

Appendix B Overview of the development and review process for this statement

i. Steps in developing and updating this statement

This statement was originally developed in November 2011 and was reviewed in July 2013 and March 2017. The Women's Health Committee carried out the following steps in reviewing this statement:

- Declarations of interest were sought from all members prior to reviewing this statement.
- The November 2011 statement was compared with the recommendations in the 10th edition of the Australian Immunisation Handbook¹ and updates were made where required.
- Recommendations were graded as set out below in Appendix B part iii).
- The updated statement was circulated out-of-session to the Women's Health Committee for comment and approval.

ii. Declaration of interest process and management

Declaring interests is essential in order to prevent any potential conflict between the private interests of members, and their duties as part of the Women’s Health Committee.

A declaration of interest form specific to guidelines and statements was developed by RANZCOG and approved by the RANZCOG Board in September 2012. The Women’s Health Committee members were required to declare their relevant interests in writing on this form prior to participating in the review of this statement.

Members were required to update their information as soon as they become aware of any changes to their interests and there was also a standing agenda item at each meeting where declarations of interest were called for and recorded as part of the meeting minutes.

There were no significant real or perceived conflicts of interest that required management during the process of updating this statement.

iii. Grading of recommendations

Each recommendation in this College statement is given an overall grade as per the table below, based on the National Health and Medical Research Council (NHMRC) Levels of Evidence and Grades of Recommendations for Developers of Guidelines. Where no robust evidence was available but there was sufficient consensus within the Women’s Health Committee, consensus-based recommendations were developed or existing ones updated and are identifiable as such. Consensus-based recommendations were agreed to by the entire committee. Good Practice Notes are highlighted throughout and provide practical guidance to facilitate implementation. These were also developed through consensus of the entire committee.

Recommendation category		Description
Evidence-based	A	Body of evidence can be trusted to guide practice
	B	Body of evidence can be trusted to guide practice in most situations
	C	Body of evidence provides some support for recommendation(s) but care should be taken in its application
	D	The body of evidence is weak and the recommendation must be applied with caution
Consensus-based		Recommendation based on clinical opinion and expertise as insufficient evidence available
Good Practice Note		Practical advice and information based on clinical opinion and expertise

Appendix C Full Disclaimer

This information is intended to provide general advice to practitioners, and should not be relied on as a substitute for proper assessment with respect to the particular circumstances of each case and the needs of any patient.

This information has been prepared having regard to general circumstances. It is the responsibility of each practitioner to have regard to the particular circumstances of each case. Clinical management should be responsive to the needs of the individual patient and the particular circumstances of each case.

This information has been prepared having regard to the information available at the time of its preparation, and each practitioner should have regard to relevant information, research or material which may have been published or become available subsequently.

Whilst the College endeavours to ensure that information is accurate and current at the time of preparation, it takes no responsibility for matters arising from changed circumstances or information or material that may have become subsequently available.